

Exercises  
**AM 0220: Nonlinear Dynamical Systems**

Bernold Fiedler, Stefan Liebscher

**due date: Mon, Feb 14, 2005**

**Problem 1:** Let  $0 < \alpha < 1$  be irrational and

$$s_n := \text{sign}(\sin(n\pi\alpha)), \quad n = 1, 2, 3, \dots$$

The sequence

$$w_n := |s_n - s_{n+1}|/2$$

detects the sign changes of the sequence  $s_n$ . Prove:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N w_n = \alpha.$$

*Free extra:* Is it possible to recover a rational number  $\alpha$  from the sequence  $s_n$ ?

**Problem 2:** Consider the vector field

$$\dot{y} = f(y), \quad y \in S^1, \quad f(y) > 0$$

with corresponding flow  $\varphi_t$ . Determine a formula for the rotation number  $\varrho(\varphi_{2\pi})$  of the time- $2\pi$ -map, for example by separation of variables.

**Problem 3:** Calculate the rotation number  $\varrho(\alpha)$  of the time- $2\pi$ -map of the differential equation

$$\dot{x} = \alpha + \sin(x - t), \quad x \in S^1.$$

**Problem 4:** Let  $f : S^1 \rightarrow S^1$  be a homeomorphism of the circle that reverses orientation.

Prove: the rotation number  $\varrho(f)$  vanishes. (Define the rotation number just as for orientation preserving homeomorphisms.)