

AM 0220: Nonlinear Dynamical Systems

Bernold Fiedler, Stefan Liebscher

due date: Mon, Apr 11, 2005**Problem 25:** Show that the system

$$\begin{aligned}\dot{x} &= x^2, \\ \dot{y} &= -y,\end{aligned}$$

has infinitely many local center manifolds of $(x, y) = (0, 0) \in \mathbb{R}^2$.*Free extra:* Modify the vector field outside a neighborhood of the origin to obtain any of these local center manifolds as a global one.**Problem 26:** Consider the system of differential equations

$$\begin{aligned}\dot{x} &= xy, \\ \dot{y} &= -y - x^2.\end{aligned}$$

Use a (local) center manifold to decide whether the equilibrium $x = y = 0$ is asymptotically stable.*Note:* Use the invariance of the center manifold to calculate the necessary terms of its Taylor expansion.**Problem 27:** Let Φ be a diffeomorphism of the plane \mathbb{R}^2 with a transverse homoclinic orbit. In class, we found a shift on two symbols for an iterate Φ^n . Prove that for every $m \in \mathbb{N}$ the shift of m symbols is conjugate to some iterate Φ^n on a suitable subset of \mathbb{R}^2 .**Problem 28:** A flow φ_t on \mathbb{R}^N is called *dissipative*, if there exists a bounded set $B \subset \mathbb{R}^N$ which absorbs every trajectory, i.e.

$$\forall x \in \mathbb{R}^N \quad \exists t_0(x) \in \mathbb{R} \quad \forall t > t_0(x) \quad \varphi_t(x) \in B.$$

Consider a dissipative flow φ_t . Prove that the set

$$A := \{x \in \mathbb{R}^N \mid \text{the orbit } \{\varphi_t(x) \mid t \in \mathbb{R}\} \text{ is bounded}\}$$

- (i) is non-empty;
- (ii) is invariant under φ_t ;
- (iii) is itself bounded;
- (iv) is attracting, i.e. $\lim_{t \rightarrow \infty} \text{dist}(\varphi_t(x), A) = 0$ for all $x \in \mathbb{R}^N$.

Free extra: The set A is an attractor, it is *the global attractor* of φ_t .